

SYSTEMIC ACQUIRED RESISTANCE

W.E. Durrant and X. Dong

*Developmental, Cell, and Molecular Biology Group, Department of Biology,
Duke University, Durham, North Carolina 27708; email: durrantw@duke.edu;
xdong@duke.edu*

Key Words plant defense, salicylic acid, SAR, NPR1, TGA factor

■ **Abstract** Systemic acquired resistance (SAR) is a mechanism of induced defense that confers long-lasting protection against a broad spectrum of microorganisms. SAR requires the signal molecule salicylic acid (SA) and is associated with accumulation of pathogenesis-related proteins, which are thought to contribute to resistance. Much progress has been made recently in elucidating the mechanism of SAR. Using the model plant *Arabidopsis*, it was discovered that the isochlorogenic acid pathway is the major source of SA during SAR. In response to SA, the positive regulator protein NPR1 moves to the nucleus where it interacts with TGA transcription factors to induce defense gene expression, thus activating SAR. Exciting new data suggest that the mobile signal for SAR might be a lipid molecule. We discuss the molecular and genetic data that have contributed to our understanding of SAR and present a model describing the sequence of events leading from initial infection to the induction of defense genes.

INTRODUCTION

Plants have evolved a number of inducible defense mechanisms against pathogen attack. Recognition of a pathogen often triggers a localized resistance reaction, known as the hypersensitive response (HR), which is characterized by rapid cell death at the site of infection (40). In the 1960s, Ross showed that tobacco plants challenged with tobacco mosaic virus (TMV) subsequently developed increased resistance to secondary infection in distal tissues (86). This spread of resistance throughout the plant's tissues was termed systemic acquired resistance (SAR). We now know that SAR can be activated in many plant species by pathogens that cause necrosis, either as part of the HR or as a symptom of disease. The resistance conferred is long-lasting, sometimes for the lifetime of the plant, and effective against a broad-spectrum of pathogens including viruses, bacteria, fungi, and oomycetes (91, 102).

Molecularly, SAR is characterized by the increased expression of a large number of pathogenesis-related genes (*PR* genes), in both local and systemic tissues. *PR* proteins were first described in the 1970s by Van Loon, who observed accumulation of various novel proteins after infection of tobacco with TMV (108, 109). Although

many PR proteins have antimicrobial properties *in vitro* (109), the function of each in the defense response has not been clearly defined. It is generally thought that SAR results from the concerted effects of many PR proteins rather than a specific PR protein. Although their roles in establishing SAR are unclear, PR genes serve as useful molecular markers for the onset of SAR.

In 1979, White observed that PR protein accumulation and resistance to TMV could be induced by treatment of tobacco with salicylic acid (SA), aspirin (acetyl SA), or benzoic acid (116). Evidence that SA is a signal for the induction of SAR came from two studies published in 1990 (63, 70). Malamy et al. showed that the endogenous SA concentration rises in both local and systemic tissues after infection of tobacco with TMV and this rise correlates with PR gene induction (63). Métraux et al. found that cucumber plants infected with either *Colletotrichum lagenarium* or tobacco necrosis virus (TNV) have considerably elevated levels of SA in the phloem sap (70). In a search for SA analogues that were less phytotoxic than SA, 2,6-dichloroisonicotinic acid (INA) and benzothiadiazole S-methyl ester (BTH) were found to induce the same set of PR genes (34, 38, 55, 69, 114). A requirement for SA as an endogenous signal for SAR was proven by Gaffney et al. using a bacterial gene, *nahG*, encoding salicylate hydroxylase, which removes SA by conversion to catechol (35). Transgenic tobacco and *Arabidopsis* expressing *nahG* accumulate very little SA after pathogen infection, fail to express PR genes, and are impaired in SAR (17, 35).

In the past 10 years, genetic analyses in the model plant *Arabidopsis* have identified additional components of SAR downstream of SA. Plants that are non-responsive to SA were identified in a number of mutant screens and found to have mutations in the same gene, *NPR1/NIM1 (NON-EXRESSER OF PR GENES1/ NONINDUCIBLE IMMUNITY1)* (8, 16, 37, 94). Considerable progress has been made in elucidating the role of NPR1 and associated proteins in the induction of SAR since the last Annual Review on SAR in 1997 (102). We therefore focus on these recent molecular and genetic experiments that have contributed to our understanding of SAR.

NATURE OF THE SYSTEMIC SIGNAL

Early grafting experiments demonstrated that the infected leaf produces a systemic signal for SAR, and this signal is not species specific (15, 46). The nature of the systemic signal has been a subject of controversy for many years.

Salicylic Acid

The detection of increased SA levels in systemic leaves and in the phloem led many researchers to believe that SA might be a systemic signal for SAR. The evidence for and against this hypothesis has been the subject of previous reviews (18, 93). Labeling studies in TMV-infected tobacco showed that most of the SA (69%) accumulating systemically was made and exported from the inoculated

leaf (97). Similarly, in cucumber infected with TNV, SA found in systemic leaves was both imported from the infected leaf and synthesized de novo (71, 73). A more recent study suggests that signaling might occur through the conversion of SA to the volatile compound methyl salicylate, which could induce resistance not only in the uninfected parts of the same plant but also in neighboring plants (98).

A number of experiments argue against SA being the systemic signal. Detachment of *Pseudomonas syringae*-infected cucumber leaves before SA levels had increased in the petiole did not block the development of SAR (85). Furthermore, grafting experiments in tobacco between wild-type scions and *nahG*-expressing rootstocks showed that, although the rootstock was unable to accumulate SA, the SAR signal was still produced and translocated to the scion (113). The reciprocal grafting experiment showed that the systemic tissue must accumulate SA for the SAR signal to be perceived.

Lipid-Based Signal Molecule

Exciting new work suggests that a lipid-based molecule may be the mobile signal for SAR. Maldonado et al. showed that the *dir1* (*defective in induced resistance 1*) mutant has normal local resistance to pathogens but is unable to develop SAR or express *PR* genes in systemic leaves (64). Therefore, wild-type DIR1, which has sequence similarity to lipid transfer proteins (LTPs), might function in the generation or transmission of the mobile signal. Indeed, experiments using petiole exudate showed that the phloem sap from *dir1* is deficient in the mobile signal for SAR. However, the mutant plants could still respond to a signal contained in the sap from wild-type plants, ruling out a role for DIR1 in signal perception. Furthermore, the *dir1* plants have wild-type SA metabolism and a normal response to SA and INA.

The similarity of DIR1 to LTPs suggests that the mobile signal for SAR might be a lipid molecule. LTPs form a multigene family in *Arabidopsis* with 71 predicted members (3). Interestingly, they share sequence similarity with elicitors from *Phytophthora* spp., which are elicitors of plant defense responses (4). The extracellular location of LTPs and elicitors is consistent with a role in signaling and implies the presence of plasma membrane (PM) receptors involved in signal transduction. Indeed, wheat LTP1 binds to the same PM receptor as the *Phytophthora* elicitor cryptogein (7).

Further evidence for a lipid-based signal molecule comes from the characterization of the *eds1* and *pad4* mutants, which are both defective in lipase-like proteins (28, 47). The *eds1* (*enhanced disease susceptibility 1*) mutant was originally identified for its compromised local resistance to *Peronospora parasitica* mediated by several resistance (*R*) genes, whereas *pad4* (*phytoalexin deficient 4*) was isolated in a screen for mutants with enhanced susceptibility to a virulent strain of *P. syringae* pv. *maculicola* (37, 81). It was subsequently discovered that *pad4* weakens local resistance mediated by the same subset of *R* genes that are blocked by *eds1* (31). These *R* genes encode TIR-NB-LRR-type resistance proteins. However,

many other *R* genes act through an EDS1-independent signaling pathway (1). In *eds1* and *pad4* plants, even when a normal HR is elicited by pathogens that trigger the EDS1-independent pathway, SAR cannot be induced (L. Jorda & J. Parker, personal communication). Experiments using phloem exudates have shown that EDS1 is required for both production of the mobile signal in the local tissue and perception of the signal in the systemic tissue (C. Lamb, personal communication). Recently, it was discovered that a tobacco SA-binding protein, SABP2 (26), is also a lipase and that its lipase activity is increased four- to fivefold by addition of SA (53). Furthermore, silencing of the *SABP2* gene diminishes both local resistance and SAR. These observations led the authors to propose that SABP2 is a receptor for SA; however, the exact position of SABP2 in the SA signaling pathway is not clear. Mutation of another gene, *SFD1*, which encodes a dihydroxyacetone phosphate reductase involved in glycerolipid synthesis, also compromises SAR and decreases SA accumulation and *PR-1* expression in systemic tissue after infection with an avirulent strain of *P. syringae* (75). Although many important questions still need to be addressed, these data strongly suggest a role for lipid signaling in SAR.

Reactive Oxygen Species

Early studies could detect no reactive oxygen species (ROS) production in systemic tissues during the onset of SAR (78, 89). However, it has since been discovered by Alvarez et al. that H_2O_2 accumulates in small groups of cells in uninoculated leaves of *Arabidopsis* after infection with an avirulent strain of *P. syringae* (2). These microbursts occur within two hours after an initial oxidative burst in the inoculated tissue and are followed by the formation of microscopic HR lesions. Using catalase to scavenge H_2O_2 , or DPI (diphenylene iodonium) to inhibit the NADPH oxidase, it was demonstrated that both the primary and secondary oxidative bursts are required for the onset of SAR. The authors propose that microbursts of ROS may activate defense responses at a low level throughout the plant and this contributes to the SAR-induced state.

Transport of the Systemic Signal

How does the SAR signal travel throughout the plant? Girdling experiments suggested that the SAR signal produced in inoculated leaves travels in the phloem to upper leaves (39, 87). If the mobile signal does travel through the phloem, the pattern of SAR induction should match the transport of sugars out of the infected leaf. When this was tested in *Arabidopsis*, it was observed that the movement of radioactively labeled sucrose did not exactly match the induction of SAR, SA accumulation, or *PR-1* expression (50). Induction of SAR was observed outside of the normal orthostichy defining phloem movement. This suggests the small amount of phloem moving between orthostichies contains enough signal to induce SAR. It seems likely that the phloem is the major conduit for the SAR signal(s), but some fraction of the signal may also be able to move by a different route.

THE ROLE OF SA IN SAR

The role of SA in SAR has been discussed extensively in a number of reviews (18, 24, 91, 93). As described above, in many plants SAR is preceded by an increase in SA concentration. However, some plants such as potato and rice have high endogenous levels of SA under noninducing conditions (14, 100, 120). Indeed, application of SA to potato does not protect it against *Phytophthora infestans* (14). However, expression of *nahG* in potato blocks resistance to *P. infestans* induced by arachidonic acid. This suggests that after treatment with arachidonic acid, instead of SA levels rising, the potato plants become more sensitive to SA (120). Thus, SA is an essential signal for SAR across a range of plants, although the mechanism by which SA induces SAR might differ.

SA Synthesis

It was previously assumed that SA for SAR is synthesized via the shikimate-phenylpropanoid pathway (57), although this was never proven. It has recently been shown that, like bacteria, plants can also synthesize SA from chorismate via isochorismate. Expression of the bacterial enzymes catalyzing these reactions, isochorismate synthase 1 (ICS1) and isochorismate pyruvate lyase 1 (IPL1), in tobacco and *Arabidopsis* results in increased SA accumulation and pathogen resistance (66, 112).

Using HPLC, Nawrath & Métraux isolated the SA induction-deficient *Arabidopsis* mutants *sid1* and *sid2*, which failed to accumulate SA after SAR induction (77). More alleles of *sid1* and *sid2*, called *eds5* and *eds16*, respectively, were identified independently by virtue of their enhanced disease-susceptibility phenotype (22, 37). A recent breakthrough in our understanding of SA biosynthesis came when *SID2/EDS16* was cloned by Wildermuth et al. and shown to encode a putative chloroplast-localized ICS1 (117). Mutations of the *ICS1* gene, in *sid2* and *eds16*, reduce SA accumulation after infection to only 5–10% of wild-type levels and compromise both basal and systemic resistance. This demonstrates that the isochorismate pathway in plants is the main source of SA synthesis during SAR. Consistent with this conclusion, *ICS1* expression is induced by infection in both local and systemic tissues. Wildermuth et al. proposed that the phenylpropanoid pathway is responsible for the rapid production of SA associated with local cell death, whereas the isochorismate pathway is more important for sustained SA synthesis during development of SAR (117). Since SA synthesis is not completely abolished in *sid2* plants, some SA must be produced either through the activity of another ICS-like protein, such as ICS2 (117), or through the phenylpropanoid pathway.

Arabidopsis ICS1 contains a putative plastid transit sequence, suggesting that SA synthesis occurs in the plastid. Interestingly, *EDS5/SID1* encodes another protein required for SA accumulation that has sequence similarity to the multidrug and toxin extrusion (MATE) family of transporter proteins (76). This suggests that

EDS5 might be involved in moving SA or a phenolic precursor out of the plastid after synthesis (68).

Control of SA Synthesis

In plants such as tobacco and *Arabidopsis*, regulation of SA biosynthesis is an essential regulatory step in SAR activation. Therefore, identification of upstream regulatory components required for the induction of SA biosynthesis genes, especially *ICS1*, will be an important step toward understanding the control of SAR. The induction of *ICS1* after infection by *Erysiphe orontii* and *P. syringae* pv. *maculicola* is not affected by depletion of SA in *nahG* plants, indicating that the *ICS1* gene is not regulated by SA (117).

Many components upstream of *ICS1* have been implicated in the regulation of SA synthesis, through characterization of various mutants with increased levels of SA. Most of these mutants form spontaneous HR-like lesions or have severe morphological phenotypes such as dwarfing (23). Expression of *ICS1* is constitutively elevated in three such gain-of-resistance mutants, *cpr1*, *cpr5*, and *cpr6* (*constitutive expresser of PR genes*) (117). However, it is unclear whether these mutants directly affect SA synthesis, or whether SA levels are elevated as an indirect effect of cell death or disruption of cellular homeostasis. It has recently been shown that two mutants with elevated SA levels, *ssi4* and *snc1*, have mutations in *R* genes that result in constitutive activation of a local defense response and therefore affect a step upstream of SA synthesis (95, 122). Interestingly, *snc1* plants do not form spontaneous lesions as observed in *ssi4*, suggesting that HR may not be required for activating SA biosynthesis.

SA synthesis induced by another *R* gene, *RPS4*, requires *EDS1* and *PAD4* (31, 125). The *eds1* and *pad4* mutants also block SA synthesis triggered by infection with virulent *P. syringae*. In *eds1* and *pad4*, induction of *EDS5*, after infection with either virulent or avirulent *P. syringae* is blocked, places *EDS1* and *PAD4* upstream of *EDS5* in the regulation of SA synthesis (76). Since *EDS1* and *PAD4* are required for resistance conferred by the same subset of *R* genes (TIR-NB-LRR) and have been shown to physically interact in planta, they are likely to function in the same pathway (31). However, the *eds1* mutation significantly impedes the onset of HR and confers full susceptibility, whereas *pad4* plants retain HR and show only intermediate susceptibility. This leads to the hypothesis that *EDS1* contributes to initial SA accumulation and development of the HR downstream of TIR-NB-LRR type *R* genes, and then recruits *PAD4* to drive amplification of the defense response by further increasing SA levels. Consistent with this hypothesis, *EDS1* and *PAD4* influence the expression of each other, with *PAD4* expression decreased more strongly in *eds1* than *EDS1* expression in *pad4* (31). This suggests that *EDS1* and *PAD4* function in a positive feedback loop that amplifies their own expression and increases production of SA after infection. A role for a positive feedback loop in SA signaling is also supported by SA-mediated *EDS1*, *PAD4*, and *EDS5* expression (28, 47, 76). The similarity of *EDS1* and *PAD4* to lipases (28, 47) suggests that lipid metabolites may be

involved in regulating the synthesis and/or accumulation of SA in local and systemic tissues.

Enhancement of the SA signal also occurs through a signal amplification loop involving ROS (96). The observation that SA binds the H₂O₂ scavenging enzymes catalase and ascorbate peroxidase (APX) and inhibits their activity led to the proposal that increases in H₂O₂ were responsible for signal transduction leading to *PR* gene induction and resistance (11, 27). However, the concentrations of SA required for inhibition of catalase and APX are higher than those seen in systemic tissues after infection. Later studies suggested that H₂O₂ functions upstream of SA. Treatment of tobacco with high concentrations (>300 mM) of H₂O₂ leads to a dose-dependent accumulation of SA and *PR-1* expression, which was suppressed in plants expressing *nahG* (58, 78). Low concentrations of SA have also been shown to potentiate the production of ROS and HR cell death. In soybean cells inoculated with *P. syringae*, the addition of SA dramatically enhances the oxidative burst and cell death (96, 103). It is hypothesized that in systemic tissues, the accumulation of low levels of SA together with the development of microbursts of ROS could amplify responses to secondary infections and contribute to SAR (25, 96).

In addition to the signal amplification loops described above, there is evidence for negative feedback of SA synthesis. In the SA-insensitive *npr1* mutant, levels of *ICS1* mRNA and SA are both elevated after infection compared to wild type (16, 94, 117). Furthermore, *npr1* mutants show reduced tolerance to exogenous SA (0.5 mM), failing to develop beyond the cotyledon stage (9, 52). The biological significance of such a feedback mechanism has yet to be determined; however, it might be utilized to shut off SAR once the pathogen challenge subsides. Many mutants with constitutively high levels of SA are dwarfs (42), and continuous spraying of wild-type plants with BTH also results in growth retardation (N. Weaver & X. Dong, unpublished observations), suggesting that accumulation of SA is detrimental to the plant's growth and development.

NPR1-DEPENDENT SA SIGNALING

To identify components involved in SA signal transduction, a number of mutant screens were performed that identified multiple alleles of a single gene, *NPR1/NIM1* (8, 16, 37, 94). Further characterization showed that the role of NPR1 is not limited to SAR. The *npr1* mutant also displays enhanced disease symptoms when infected with virulent pathogens and is impaired in some *R* gene-mediated resistance, suggesting that NPR1 is important for restricting the growth of pathogens at the site of infection (8, 16, 37, 94). NPR1 is required for another induced resistance response, known as induced systemic resistance (ISR), which is triggered by nonpathogenic root-colonizing bacteria and confers resistance to bacteria and fungi in aerial parts of the plant (82, 83). NPR1 also mediates cross-talk between the SA signaling pathway and the jasmonic acid (JA) and ethylene (C₂H₄) signaling pathways that confer resistance to insects and some necrotrophic pathogens (101). As discussed earlier, *npr1* has reduced tolerance to SA toxicity and accumulates

high endogenous levels of SA, suggesting a role for NPR1 in both detoxification of SA and feedback regulation of SA biosynthesis (9, 52). In addition, NPR1 has functions that are not directly related to resistance, including the regulation of cell division and/or endoreduplication (111).

NPR1 is expressed throughout the plant at low levels and its mRNA levels rise two- to threefold after pathogen infection or treatment with SA (9, 90). *NPR1* expression is likely mediated by WRKY transcription factors as mutation of the WRKY binding sites (W-boxes) in the *NPR1* promoter abolished its expression (119). Overexpression of *NPR1* in *Arabidopsis* enhances resistance to *P. parasitica*, *P. syringae*, and *Erysiphe cichoracearum* with no apparent detrimental effects on the plant (10, 33). Unlike many mutants with constitutive resistance, the *NPR1* overexpressing lines do not constitutively express *PR* genes. The enhancement of resistance is probably caused by the more rapid or higher induction of *PR* genes observed in these overexpressing lines (10, 33). This indicates that, even when expressed at higher levels, the NPR1 protein must be activated to induce SAR.

The NPR1 protein has two protein-protein interaction domains, an ankyrin-repeat and a BTB/POZ (*Broad-Complex, Tramtrack, Bric-a-brac/Poxvirus, Zinc finger*) domain, as well as a putative nuclear localization signal and phosphorylation sites (9, 90). Functional studies have shown that accumulation of NPR1 in the nucleus after treatment with SAR inducers is essential for *PR* gene induction (52).

NPR1 homologues have been identified in rice, tobacco, tomato, apple, and orange (12, 61; S.-Y. He, personal communication; M. Kinkema, J.-Y. Yang & X. Dong, unpublished observations), suggesting that NPR1 function is conserved across plant species. This is supported by the demonstration that overexpression of *Arabidopsis NPR1* in rice confers resistance to *Xanthomonas oryzae* pv. *oryzae* (12).

TGA Transcription Factors

The absence of any obvious DNA-binding domain and the presence of protein-protein interaction domains in NPR1 prompted several laboratories to carry out yeast two-hybrid screens for NPR1-interacting proteins. In one of these screens, three small structurally similar proteins named NIMIN1, NIMIN2, and NIMIN3 (NIM interactor) were identified. NIMIN1 and NIMIN2 interact with the C terminus of NPR1, while NIMIN3 interacts with the N terminus (115). NIMINs contain stretches of acidic amino acids and are hypothesized to be transcription factors; however, more experiments are required to demonstrate their biological activity.

The predominant NPR1 interactors found in the yeast two-hybrid screens were members of the TGA family of basic leucine zipper transcription factors. NPR1 interacts with the *Arabidopsis* TGA factors, TGA2, TGA3, TGA5, TGA6, and TGA7 but only weakly or not at all with TGA1 and TGA4 (20, 51, 121, 124). NPR1 also interacts with TGA factors from tobacco and rice (12, 80). Using truncated or mutant forms of NPR1, the ankyrin-repeat domain in the middle of the protein

was shown to be essential for binding TGA factors, while the N-terminal region appears to enhance binding (20, 121, 124).

TGA factors bind to activator sequence-1 (*as-1*) or *as-1*-like promoter elements (49), which have been found in several plant promoters activated during defense, including *Arabidopsis PR-1* (56). Linker scanning mutagenesis of the *PR-1* promoter identified two *as-1*-like elements, *LS7* and *LS5*. *LS7* is a positive regulatory element required for induction by INA, whereas *LS5* is a weak negative regulatory element (56). Després et al. used these *cis*-elements as probes for electrophoretic mobility shift assays (EMSA) and showed that both TGA2 and TGA4 could bind to *LS7*, whereas only TGA2 could bind to *LS5* (20). Furthermore, binding of TGA2 but not TGA4 was enhanced by the addition of NPR1, consistent with the yeast two-hybrid interaction data.

Although NPR1 is clearly a positive regulator of *PR* genes, it may exert its function by either enhancing a transcriptional activator or inhibiting a transcriptional repressor. The presence of multiple *as-1*-like elements in the *PR-1* promoter and the differential binding affinities of each TGA factor to these elements as well as to NPR1 highlight the complexity of the regulatory mechanism. Indeed, in an EMSA performed by Després et al., binding to the *as-1* element from the *35S* promoter was significantly enhanced in protein extracts from SA-treated plants (20). However, extracts from untreated *npr1* plants also contained strong *as-1*-binding activity and this was not changed by SA treatment. This result can be reconciled if different TGA factors are responsible for the observed *as-1*-binding in wild type and *npr1*. Consistent with this hypothesis, when EMSA was performed using the *LS7* element from the *PR-1* promoter, Johnson et al. found that SA enhanced the binding activity in wild-type plants and this binding was abolished in *npr1* plants (48). This result was confirmed by chromatin immunoprecipitation (ChIP) experiments (48). Recruitment of TGA2 or TGA3 to a 1kb fragment of the *PR-1* promoter was observed only after treatment with SA and not in untreated wild-type plants or SA-treated *npr1* plants. Unfortunately, ChIP is unable to resolve the adjacent binding sites *LS7* and *LS5*, which are within 30 bp of each other.

Another approach to study the role of TGA factors in vivo is to examine the phenotypes of mutant plants. As there are 10 TGA factors in *Arabidopsis* (45), functional redundancy may prevent observation of a mutant phenotype. Indeed, analysis of single knockout mutants of TGA2 and TGA3 revealed little phenotype (M. Kesarwani & X. Dong, unpublished observations). Consistent with this, overexpression or silencing of TGA2 did not alter resistance to a virulent strain of *P. parasitica* (51). However, overexpression of TGA5 enhanced resistance to *P. parasitica*, but this was not dependent on SA or NPR1 and did not correlate with *PR* gene expression. Using a reverse genetics approach, Li et al. isolated a knockout of the adjacent *TGA2* and *TGA5* genes (59). This was crossed to a knockout of *TGA6* to create the *tga2 tga5 tga6* triple mutant (123), thus deleting all members of one of three subclasses of TGA factors (118). The *tga2 tga5 tga6* triple mutant has phenotypes similar to *npr1*, showing compromised SAR and decreased tolerance to high concentrations of SA. All three genes must be deleted to

observe this phenotype, leading to the conclusion that TGA2, TGA5, and TGA6 are essential for and play redundant roles in the induction of SAR. Interestingly, the triple knockout and also the *tga2 tga5* double mutant have increased *PR-1* expression in the absence of SAR induction, suggesting that TGA factors also play a role in the repression of basal *PR-1* expression. This might be through interaction with the negative *LS5* element in the *PR-1* promoter (56). If knocking out *TGA2*, *TGA5*, and *TGA6* is sufficient to abolish SAR, then what is the role of the other TGA factors? Despite compromised SAR, the *tga2 tga5 tga6* triple mutant does not show enhanced susceptibility to a virulent strain of *P. syringae*, suggesting that other TGA factors may be involved in basal resistance. It is also possible that the different subgroups of TGA factors regulate different sets of defense genes and that loss of induction of one set would be sufficient to impede SAR. This question can best be addressed by deleting all members of the other subgroups of TGA factors.

As an alternative to mutant analysis, dominant-negative versions of TGA factors that can no longer bind to DNA were expressed in tobacco and *Arabidopsis*. In tobacco, overexpression of a dominant-negative *TGA2.2* decreased *as-1*-binding activity and *PR* gene induction (79). In another study, a dominant-negative version of *Arabidopsis* *TGA2* was expressed in tobacco (84). In this case, *as-1*-binding activity was completely abolished in untreated plants but in contrast to the previous study, induction of *PR* genes and pathogen resistance were both enhanced. The discrepancy between these experiments is difficult to explain because the dominant-negative TGA factors used in these studies could bind endogenous TGA factors as well as NPR1. However, these studies do support the idea that TGA factors can play both positive and negative roles in *PR* gene regulation.

To observe activity of specific TGA factors *in vivo*, chimeric transcription factors have been constructed in which *TGA2* or *TGA3* were fused to the yeast GAL4 DNA-binding domain. Fan & Dong showed that replacing the bZIP domain of *TGA2* with the GAL4 DNA-binding domain produced a transcription factor that activated the expression of a *UAS^{GAL4}::GUS* reporter construct in response to INA or SA (29). This reporter gene activation was abolished in the *npr1* mutant. EMSA showed that *TGA2*-GAL4 binding to *UAS^{GAL4}* was enhanced by INA treatment and is dependent on NPR1. This is consistent with the findings that NPR1 enhances binding of *TGA2* to the *LS7* element (20). Johnson et al. used a similar heterologous system to show that *TGA3* is also a transcriptional activator (48).

Redox Signaling

The *in vivo* interaction of NPR1 with TGA factors requires induction with SA, even though both proteins are constitutively expressed. Until recently, the controlling mechanisms for NPR1 nuclear localization and activation of TGA factors were unclear. Two exciting new papers have revealed that changes in the redox status of the cell after SA treatment play an important role in this regulation (19, 74).

The observation that NPR1-like proteins from different species contain ten conserved cysteines suggested that NPR1 might be under redox-regulation. Mou et al. tested this hypothesis by examining NPR1 under different redox conditions (74). When proteins were extracted in the absence of the reducing agent DTT, monomeric NPR1 could only be detected in INA-treated samples, whereas in the presence of DTT equal amounts of monomeric NPR1 were detected with or without INA treatment. This suggests that before and after SAR induction NPR1 is present in two different conformations. As antibodies developed against the endogenous NPR1 only recognize the reduced form, further analyses used GFP-tagged NPR1. In extracts from untreated plants, NPR1-GFP was found in a high MW (>250,000) protein complex. Addition of DTT to the sample reduced all the NPR1-GFP to its monomeric form, suggesting that the high MW complex is formed through intermolecular disulfide bridges. Interestingly, this complex is partially reduced as a result of INA treatment. This pattern of conformational changes was also observed after pathogen infection in both inoculated and distal tissues and precedes *PR* gene induction. Measurements of cellular glutathione pools showed that a biphasic change in redox occurs after SAR induction, first oxidizing and then reducing. Lowering NADPH levels diminished both NPR1-GFP reduction and *PR-1* expression after INA treatment. Mutation of cysteine residues C82 or C216 in NPR1 resulted in constitutive monomerization, nuclear accumulation of mutant proteins and expression of *PR-1*. These data demonstrate that the monomer is the active form of NPR1 for *PR-1* induction and provide a model for the activation of NPR1 by SA: SA accumulation triggers conversion of NPR1 from an oligomer to a monomer through changes in cellular redox status favoring reduction. This monomeric form of NPR1 is then able to move to the nucleus where it interacts with TGA factors to induce *PR* gene expression.

As discussed earlier, in yeast two-hybrid studies NPR1 interacts strongly with TGA2 and TGA3 but very weakly or not at all with TGA1 and TGA4 (20, 124). Using a plant two-hybrid assay in *Arabidopsis*, Després et al. demonstrated a physical interaction between NPR1 and TGA1 (19). Using domain swapping between TGA1 and TGA2, the plant-specific regulatory region was defined to a 30 aa region containing two cysteine residues in TGA1 (and TGA4) that are not found in TGA2 or other TGA factors. Mutation of these residues in TGA1 allowed interaction with NPR1 in yeast and in untreated leaves. A clever labeling experiment designed to distinguish between reduced and oxidized cysteine residues showed that TGA1 (and/or TGA4) exists in both oxidized and reduced forms in untreated leaves. After SA treatment, only the reduced form was detected (19). This indicates that SA controls the redox status of TGA1 (and/or TGA4) and only the reduced form can bind NPR1. Evidently, redox changes do not increase TGA1 DNA-binding activity directly; instead, this is achieved through enhancement of its interaction with NPR1. These experiments add TGA1 and TGA4 to the spectrum of TGA factors that interact with NPR1, although their role in SAR has yet to be established. These results are consistent with the finding by Mou et al. that the SA signal is transduced through changes in cellular reduction potential that lead to

monomerization of NPR1 (74). We are just beginning to realize the significance of such redox changes during SAR. In the past, research has mainly focused on the initial oxidative burst during *R* gene-mediated defense. These two studies highlight the importance of the reducing state that usually appears following the oxidative stress.

OTHER REGULATORY COMPONENTS IN SAR SIGNALING

To identify other components of SAR signaling, several genetic screens have been conducted looking for suppressors of *npr1*. The recessive *snl1* (*suppressor of npr1 inducible*) mutant restores SA-inducible *PR* gene expression and pathogen resistance in the *npr1* background (60). The *snl1* mutant has wild-type levels of SA and only slightly elevated expression of *PR* genes in the absence of an SAR inducer, indicating that *snl1* is likely downstream of *npr1*. The *snl1* phenotype suggests that SNI1 is a negative regulator of *PR* gene expression and SAR, whose repression is relieved by NPR1 after induction. The low basal levels of *PR* gene expression and restored induction in the *snl1 npr1* double mutant indicate that, in addition to NPR1-mediated inactivation of SNI1, an SA-dependent but NPR1-independent regulatory step is also required for SAR gene induction. Consistent with the hypothesis that SNI1 is a repressor of *PR* genes, SNI1-GFP has been observed in the nucleus when bombarded into onion epidermal cells (60). Moreover, when fused to the GAL4 DNA-binding domain and expressed in yeast, SNI1 repressed transcription of a reporter carrying *UAS^{GAL4}* upstream of a constitutive promoter (R. Mosher & X. Dong, unpublished observations). This result suggests that SNI1 may repress a general transcriptional mechanism that is conserved between yeast and plants.

SNI1 is a novel plant-specific protein with no similarity to proteins of known function. However, putative homologues have been found in many plant species including barley, *Medicago truncatula*, potato, rice, soybean, and sugar cane, indicating that SNI1 function may be conserved (R. Mosher & X. Dong, unpublished observations). SNI1 contains no obvious DNA-binding domain, suggesting that it represses transcription through interaction with other factors rather than binding directly to the promoter. Linker scanning mutagenesis of the *PR-1* promoter identified the negative regulatory *cis*-element, *LS4*, which has a W-box consensus sequence (56). Mutation of *LS4* resulted in elevated basal expression and stronger induction in response to INA treatment, a pattern similar to that of *PR-1* in *snl1*. This suggests that SNI1 might be recruited to the *PR-1* promoter through interaction with a WRKY factor.

As an alternative approach to investigate the function of SNI1, a screen for suppressors of *snl1* (*ssn*) was performed. Three mutants were identified, *ssn1*, *ssn2*, and *ssn3*, which alleviate both the dwarf morphology and the background *PR* gene expression of *snl1* (W. Durrant & X. Dong, unpublished observations). In the *snl1 ssn* double mutants, the pattern of *PR* gene expression is the same

as wild-type, whereas in the *snl npr1 ssn* triple mutants induction of *PR* gene expression by SA is blocked, similar to *npr1*. These results indicate that the wild-type *SSN* genes are involved in controlling both basal and SA-inducible *PR* gene expression observed in *snl npr1*. In other words, the same transcription factor(s) is probably responsible for the background and SA-inducible NPR1-independent *PR* gene expression.

Likely candidates for regulators of NPR1-independent *PR* gene expression and resistance are the Whirly (Why) family of transcription factors, named after the whirling appearance of their crystal structure (21). Potato StWhy1 specifically binds the single-stranded form of a *cis*-element in the *PR10a* promoter. *Arabidopsis* has three genes encoding Whirly proteins, of which AtWhy1 is the most similar to StWhy1 (21a). SA treatment induced AtWhy1 DNA-binding activity in both wild-type and *npr1* plants, indicating that AtWhy1 activation is independent of NPR1. Knockout mutants of *AtWhy1* were lethal. However, two lines with mis-sense mutations in the ssDNA-binding domain (*atwhy1.1*) or the central α -helical region (*atwhy1.2*) were viable and showed reduced DNA-binding activity, SA-induced *PR-1* transcript accumulation, and resistance to *P. parasitica*. The NPR1-independent activation of AtWhy1 and the decrease in *PR-1* expression observed in the *atwhy1* mutants suggest that AtWhy1 is important for NPR1-independent *PR* gene expression. It would be interesting to test whether *atwhy1* mutants can block the NPR1-independent *PR* gene expression observed in *snl npr1*.

Other possible SAR regulatory components include DTH9 (DETACHMENT 9). The *dth9* mutant has increased susceptibility to virulent pathogens, accumulates elevated levels of SA, and fails to develop SAR in response to pathogen infection or SA treatment (67). These phenotypes are reminiscent of *npr1*; however, *dth9* differs from *npr1* in that its *PR-1* and *PR-2* expression in response to infection or SA treatment is unaltered. Since SA treatment did not reverse the disease susceptibility observed in the mutant, DTH9 should be placed downstream of SA in a pathway parallel to NPR1 that contributes to SAR. Interestingly, *dth9* is also insensitive to auxin treatment, indicating that auxin signaling may play a role in plant defense.

GENE EXPRESSION CHANGES DURING SAR

There is ample evidence indicating that SAR is conferred by expression of a collection of genes. The phenotype of *dth9* is a good example, showing that SAR can be blocked without affecting the induction of *PR-1* and *PR-2*. The sequencing of the *Arabidopsis* genome has allowed global analyses of gene expression changes during SAR to be conducted using DNA microarray technology. Maleck et al. surveyed 25–30% of *Arabidopsis* genes under 14 SAR-inducing or repressing conditions, identifying 413 ESTs (about 300 genes) that show differential expression during SAR (65). Cluster analysis was used to identify a group of 31 genes with a similar transcription pattern to that of *PR-1*. The genes in this *PR-1* regulon are strongly

induced in systemic tissue during SAR and this induction is NPR1-dependent. They are also induced by infection with virulent *P. parasitica*, suggesting that activation of SAR-related genes in local tissue can limit infection by compatible pathogens. This is consistent with the enhanced susceptibility to virulent pathogens observed in *npr1* mutants. Interestingly, only 17 of 26 *PR-1* regulon genes have an *as-1* element in their promoter, whereas W-boxes were overrepresented, occurring an average of 4.3 times in every promoter. This suggests that WRKY factors rather than TGA factors are important for coregulation of *PR-1* regulon genes. The *LS4* W-box in the *PR-1* promoter acts as a strong negative *cis*-element (56), leading Maleck et al. to propose that WRKY factors repress the expression of *PR-1* regulon genes (65). Upon activation of SAR, NPR1-dependent derepression would occur, possibly through the inactivation of SNII.

Promoter analysis was also performed in another study on 1058 genes that were induced by pathogen infection, SA, MeJA, or ozone (62). This revealed that *as-1* elements, W-boxes, abscisic acid response elements, and G-boxes were overrepresented across all treatments, whereas the Myc motif was overrepresented only in the SA-induced genes. This suggests a role for these *cis*-elements in stress responses but does not identify which are important during SAR.

A different use of microarray analysis is to identify the global expression phenotype of mutants impaired in disease resistance (36). This allowed the placement of three mutants with previously undefined roles into known signal pathways: *pad1* and *eds8* in the JA/C₂H₄ pathway and *eds3* in the SA pathway. This result was confirmed by the demonstration that *pad1* and *eds8* are impaired in JA-induced anthocyanin production and *eds3* has greatly reduced SA production after infection with a virulent strain of *P. syringae*. This study also revealed that two mutant alleles of *NPR1* have different effects on pathogen-induced gene expression. The *npr1-3* mutant appears to affect only expression of SA-regulated genes, whereas *npr1-1* also affects expression of genes that require JA and C₂H₄ for induction. This may indicate that the C-terminal 194 aa of NPR1, which are missing in *npr1-3* (9), are not required for JA/C₂H₄-induced gene expression. It will be interesting to test whether *npr1-3* is impaired in ISR, which involves JA/C₂H₄-signaling instead of SA signaling. The mutation in *npr1-1* changes a structurally important residue in the ankyrin-repeat domain and likely results in a misfolded, nonfunctional protein. In *npr1-1* both SAR and ISR are blocked (8, 83).

A potential difficulty in interpreting microarray data is the lack of hierarchical information on the transcriptional events that occur during SAR. The SAR-induced genes include effector genes that confer resistance as well as regulatory genes such as transcription factors. To overcome this problem, a strategy was devised to focus on only a single transcriptional level (D. Wang & X. Dong, unpublished data). Fusion of NPR1 to the hormone-binding domain of the rat glucocorticoid receptor (GR) placed its nuclear localization under the control of the steroid hormone dexamethasone (52). Treating an *NPR1-GR* transgenic line (in *npr1* background) with SA will activate expression of SA-mediated NPR1-independent genes. Addition of dexamethasone after SA treatment will allow NPR1-GR to move to the nucleus and induce NPR1-mediated genes. To focus on the direct transcriptional targets

of NPR1, cycloheximide can be added to inhibit de novo protein synthesis and secondary transcriptional events.

INTERACTION BETWEEN SAR AND OTHER DEFENSE PATHWAYS

It is impossible to understand SAR fully without studying its interaction with other biological processes. It is hypothesized that plant defense pathways interact synergistically or antagonistically to fine-tune responses according to the challenging organism(s). Different responses may confer resistance to the same pathogen. On the other hand, activation of one pathway may lead to cross-talk inhibition of another that is less effective against the pathogen. Cross-talk between different defense pathways is reviewed elsewhere (5). Here, we focus on interactions involving components of SAR.

As mentioned earlier, NPR1 is required for other induced resistance pathways, including ISR induced by the nonpathogenic bacterium *P. fluorescens* (83, 107). Like SAR, ISR protects plants against a range of pathogens (106), but it is independent of SA and PR gene induction (82). ISR is blocked in the *jar1* (*jasmonic acid resistant 1*) and *etr1* (*ethylene resistant 1*) mutants, indicating a requirement for JA and C₂H₄ signaling components (83). The *Arabidopsis* ecotypes RLD and Ws, which fail to develop ISR, carry a recessive mutation that causes C₂H₄-insensitivity (104). Two previously identified mutants, *eds4* and *eds8*, which are insensitive to C₂H₄ and JA, respectively, are also impaired in ISR (105). However, a third *eds* mutant, *eds10*, responds normally to both hormones and develops normal SAR but is blocked for ISR, suggesting that *EDS10* is a novel component of ISR. ISR requires NPR1 at a point downstream of JAR1 and ETR1 (83). NPR1 is therefore an important regulator of induced defense responses downstream of either SA or JA/C₂H₄ and may differentially regulate these responses according to upstream signals. Furthermore, simultaneous induction of SAR and ISR has an additive effect on the level of induced resistance against *P. syringae* (110). Therefore, NPR1 is able to function in both of these pathways simultaneously.

There are also examples of cross-resistance where insect feeding can induce aspects of SAR (43). This has been observed in response to aphids and whiteflies, which are sucking insects and therefore do minimal damage to the tissue. The idea that plants perceive some insects as pathogens rather than herbivores is supported by the identification of an *R* gene that confers resistance to aphids and nematodes (72, 88). It makes sense for the plant to activate SAR when attacked by such insects if they act as vectors for pathogens. Evidence for coregulation by SA and JA signaling also comes from a gene expression profiling study in which 55 genes were induced by treatment with either SA or JA (92).

Besides synergy described above, there is increasing evidence for antagonism between the SA and JA/C₂H₄ pathways (43, 54). For example, tobacco plants with enhanced SAR have decreased systemic resistance to the chewing insect *Heliothis virescens* after SAR induction, whereas plants with reduced SAR show

more effective resistance (30). The induction of SAR therefore has a negative effect on the JA/C₂H₄ pathways, which are normally induced by chewing insects and wounding. SA accumulation has been associated with inhibition of JA biosynthesis and decreased expression of JA-responsive genes (43). A large-scale analysis of gene expression in wild-type and mutants defective in SA, JA or C₂H₄ signaling revealed that, although JA/C₂H₄ signaling can sometimes inhibit SA signaling, repression of JA signaling by SA signaling is more prevalent (36). Treatment of *Arabidopsis* with SA and JA simultaneously prevents the expression of JA-responsive genes (101). Studies with *npr1* plants showed that this antagonistic effect of SA on JA signaling requires NPR1. Unlike the activation of SAR, nuclear localization of NPR1 was not required for suppression of JA signaling. Thus, NPR1 is a central regulator of plant defense responses including SAR, ISR, and SA/JA cross-talk. A challenge for the future will be to understand how NPR1 coordinates these different responses and to unravel the signaling network downstream of NPR1 in each case.

FITNESS COSTS OF SAR

It has often been suggested that disease resistance is associated with fitness costs and that plants have evolved inducible defense mechanisms because it is too costly to have defense responses switched on all the time (6, 41, 42). The phenotypes of many mutants that show constitutive *PR* gene expression, accumulation of SA, and resistance to pathogens support this idea. These mutants often have reduced plant size, loss of apical dominance, curly leaves, and decreased fertility, all traits associated with decreased plant fitness (reviewed in 42). Consistent with this view, overexpression of NPR1 in rice triggers lesion development and chlorosis under certain environmental conditions, correlating with expression of defense genes (32). Although only a few studies have specifically addressed the fitness costs of SAR, they all conclude that constitutive expression of SAR in uninfected plants is detrimental.

The SA analogue BTH, which has been developed commercially as a plant activator, induces SAR in wheat and confers systemic protection against powdery mildew (38). Heil et al. have therefore studied the fitness costs associated with the use of BTH in the absence of pathogens (44). Plants grown hydroponically in the absence of pathogens or in the field showed a reduction in biomass and the number of ears and grains if treated with BTH. These effects were stronger when combined with limited nitrogen availability. Similarly, treatment of *Arabidopsis* with SA decreases seed yield (13). These experiments, although not representative of natural conditions, are useful when considering the costs and benefits of treating crop plants with such compounds.

An alternative to exogenous chemical treatment is to examine the fitness of mutants with constitutive SAR or mutants impaired in SAR. Two studies have shown that gain-of-resistance mutants such as *cpr1*, *cpr5*, *cpr6*, and *cep1* [*constitutive expression of PR-1*; (99)] have decreased fitness, demonstrated by low seed yield and small rosette diameter (13; A. Heidel, J. Antonovics & X. Dong, unpublished




Figure 1 The sequence of events from pathogen recognition to defense gene induction.

observations). The cost of resistance observed here could be due to the allocation of the plant's resources to constitutive PR protein production. As well as a cost of constitutive SAR, there is a fitness cost associated with the inability to induce SAR. One study showed that *nahG* and *nim1* plants grown in a growth chamber have high seed production in the absence of SAR induction (13). However, in a field experiment, the *npr1* mutant and the *NPR1-L* transgenic line, in which *NPR1* is silenced (10), showed a reduction of fitness (A. Heidel, J. Antonovics & X. Dong, unpublished observations). Although there was no visible sign of disease on these plants, they might have had a low level of infection that was detrimental to plant growth. It is also possible that soil bacteria could have induced ISR in the wild-type plants, but not in *npr1* or the *NPR1-L* line, to protect them from microbial pathogens. This study shows that SAR has fitness benefits and fitness costs consistent with the inducible nature of this defense response. Such results

have important implications for development of crop protection strategies through manipulation of SAR.

CONCLUSIONS

Our understanding of SAR has increased considerably over recent years as we have begun to elucidate the molecular mechanisms underlying this response. In Figure 1, we present a summary of the data discussed in this review. Many of the processes contributing to SAR are clearly required in both local and systemic tissues and contribute to basal disease resistance. These include the synthesis of SA, changes in redox status, and the induction of defense gene expression. In local tissue, the trigger for these changes is the recognition of the invading pathogen, whereas in systemic tissue they are induced by perception of a systemic signal. There is evidence for negative and positive feedback of SA signaling and cross-talk between different signaling pathways, adding to the complexity of the defense response. As well as the central role played by NPR1-mediated signaling, there is growing evidence for an NPR1-independent pathway(s) that contributes to defense gene induction. Challenges for the future include identification of the mobile signal for SAR, to which we are one step closer after the identification of DIR1. Induction of SAR to control infection of crop plants is already being used in the field by application of BTH and it has been suggested that NPR1 overexpression is another viable strategy. Better understanding of the SAR signaling pathway will certainly lead to new environmentally friendly methods of crop protection.

ACKNOWLEDGMENTS

We thank Lucia Jorda, Jane Parker, Chris Lamb, Sheng-Yang He, Darrell Desveaux, Rajagopal Subramaniam, Jeff Dangl, Normand Brisson, and our colleagues in the Dong lab for sharing unpublished data. We thank Lisa Anderson, Meenu Kesarwani, Rebecca Mosher, Zhonglin Mou, Graham Redman, and Natalie Weaver for critical reading of the manuscript. W. Durrant is supported by a fellowship from The International Human Frontier Science Program Organization. X. Dong is supported by grants from United States Department of Agriculture and National Science Foundation.

The *Annual Review of Phytopathology* is online at <http://phyto.annualreviews.org>

LITERATURE CITED

1. Aarts N, Metz M, Holub E, Staskawicz BJ, Daniels MJ, Parker JE. 1998. Different requirements for *EDS1* and *NDR1* by disease resistance genes define at least two *R* gene-mediated signalling pathways in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* 95:10306–11
2. Alvarez ME, Pennell RI, Meijer P-J,

- Ishikawa A, Dixon RA, Lamb C. 1998. Reactive oxygen intermediates mediate a systemic signal network in the establishment of plant immunity. *Cell* 92:773–84
3. Beisson F, Koo AJ, Ruuska S, Schwender J, Pollard M, et al. 2003. Arabidopsis genes involved in acyl lipid metabolism. A 2003 census of the candidates, a study of the distribution of expressed sequence tags in organs, and a web-based database. *Plant Physiol.* 132:681–97
 4. Blein J-P, Coutos-Thévenot P, Marion D, Ponchet M. 2002. From elicitors to lipid-transfer proteins: a new insight in cell signalling involved in plant defence mechanisms. *Trends Plant Sci.* 7:293–96
 5. Bostock RM. 1999. Signal conflicts and synergies in induced resistance to multiple attackers. *Physiol. Mol. Plant Pathol.* 55:99–109
 6. Brown JKM. 2002. Yield penalties of disease resistance in crops. *Curr. Opin. Plant Biol.* 5:339–44
 7. Buhot N, Douliez J-P, Jacquemard A, Marion D, Tran V, et al. 2001. A lipid transfer protein binds to a receptor involved in the control of plant defence responses. *FEBS Lett.* 509:27–30
 8. Cao H, Bowling SA, Gordon S, Dong X. 1994. Characterization of an Arabidopsis mutant that is nonresponsive to inducers of systemic acquired resistance. *Plant Cell* 6:1583–92
 9. Cao H, Glazebrook J, Clark JD, Volko S, Dong X. 1997. The Arabidopsis *NPR1* gene that controls systemic acquired resistance encodes a novel protein containing ankyrin repeats. *Cell* 88:57–63
 10. Cao H, Li X, Dong X. 1998. Generation of broad-spectrum disease resistance by overexpression of an essential regulatory gene in systemic acquired resistance. *Proc. Natl. Acad. Sci. USA* 95:6531–36
 11. Chen Z, Silva H, Klessig DF. 1993. Active oxygen species in the induction of plant systemic acquired resistance by salicylic acid. *Science* 262:1883–86
 12. Chern M-S, Fitzgerald HA, Yadav RC, Canlas PE, Dong X, Ronald PC. 2001. Evidence for a disease-resistance pathway in rice similar to the *NPR1*-mediated signaling pathway in *Arabidopsis*. *Plant J.* 27:101–13
 13. Cipollini DF. 2002. Does competition magnify the fitness costs of induced responses in *Arabidopsis thaliana*? A manipulative approach. *Oecologia* 131:514–20
 14. Coquoz J-L, Buchala A, MP H, Métraux JP. 1995. Arachidonic acid induces local but not systemic synthesis of salicylic acid and confers systemic resistance in potato plants to *Phytophthora infestans* and *Alternaria solani*. *Phytopathology* 85:1219–24
 15. Dean RA, Kuc J. 1986. Induced systemic protection in cucumbers: the source of the “signal”. *Physiol. Mol. Plant Pathol.* 28:227–33
 16. Delaney TP, Friedrich L, Ryals JA. 1995. *Arabidopsis* signal transduction mutant defective in chemically and biologically induced disease resistance. *Proc. Natl. Acad. Sci. USA* 92:6602–6
 17. Delaney TP, Uknes S, Vernooij B, Friedrich L, Weymann K, et al. 1994. A central role of salicylic acid in plant disease resistance. *Science* 266:1247–50
 18. Dempsey DA, Shah J, Klessig DF. 1999. Salicylic acid and disease resistance in plants. *Crit. Rev. Plant Sci.* 18:547–75
 19. Després C, Chubak C, Rochon A, Clark R, Bethune T, et al. 2003. The Arabidopsis *NPR1* disease resistance protein is a novel cofactor that confers redox regulation of DNA binding activity to the basic domain/leucine zipper transcription factor TGA1. *Plant Cell* 15:2181–91
 20. Després C, DeLong C, Glaze S, Liu E, Fobert PR. 2000. The Arabidopsis *NPR1/NIM1* protein enhances the DNA binding activity of a subgroup of the TGA family of bZIP transcription factors. *Plant Cell* 12:279–90
 21. Desveaux D, Allard J, Brisson N,

- Sygusch J. 2002. A new family of plant transcription factors displays a novel ssDNA-binding surface. *Nat. Struct. Biol.* 9:512–17
- 21a. Desveaux D, Subramaniam R, Després C, Mess J-N, Lévesque C, et al. 2004. A “Whirly” transcription factor is required for salicylic acid-dependent disease resistance in *Arabidopsis*. *Dev. Cell* 6:229–40
 22. Dewdney J, Reuber TL, Wildermuth MC, Devoto A, Cui J, et al. 2000. Three unique mutants of *Arabidopsis* identify *eds* loci required for limiting growth of a biotrophic fungal pathogen. *Plant J.* 24:205–18
 23. Dietrich RA, Delaney TP, Uknes SJ, Ward ER, Ryals JA, Dangel JL. 1994. *Arabidopsis* mutants simulating disease resistance response. *Cell* 77:565–77
 24. Dong X. 2001. Genetic dissection of systemic acquired resistance. *Curr. Opin. Plant Biol.* 4:309–14
 25. Draper J. 1997. Salicylate, superoxide synthesis and cell suicide in plant defence. *Trends Plant Sci.* 2:162–65
 26. Du H, Klessig DF. 1997. Identification of a soluble, high-affinity salicylic acid-binding protein in tobacco. *Plant Physiol.* 113:1319–27
 27. Durner J, Klessig DF. 1995. Inhibition of ascorbate peroxidase by salicylic acid and 2,6-dichloroisonicotinic acid, two inducers of plant defense responses. *Proc. Natl. Acad. Sci. USA* 92:11312–16
 28. Falk A, Feys BJ, Frost LN, Jones JDG, Daniels MJ, Parker JE. 1999. *EDS1*, an essential component of *R* gene-mediated disease resistance in *Arabidopsis* has homology to eukaryotic lipases. *Proc. Natl. Acad. Sci. USA* 96:3292–97
 29. Fan W, Dong X. 2002. In vivo interaction between NPR1 and transcription factor TGA2 leads to salicylic acid-mediated gene activation in *Arabidopsis*. *Plant Cell* 14:1377–89
 30. Felton GW, Korth KL, Bi JL, Wesley SV, Huhman DV, et al. 1999. Inverse relationship between systemic resistance of plants to microorganisms and to insect herbivory. *Curr. Biol.* 9:317–20
 31. Feys BJ, Moisan LJ, Newman M-A, Parker JE. 2001. Direct interaction between the *Arabidopsis* disease resistance signaling proteins, EDS1 and PAD4. *EMBO J.* 20:5400–11
 32. Fitzgerald HA, Chern M-S, Navarre R, Ronald PC. 2004. Over-expression of (*At*)*NPR1* in rice leads to a BTH- and environment-inducible lesion-mimic/cell death phenotype. *Mol. Plant-Microbe Interact.* 17:140–51
 33. Friedrich L, Lawton K, Dietrich R, Willits M, Cade R, Ryals J. 2001. *NIMI* overexpression in *Arabidopsis* potentiates plant disease resistance and results in enhanced effectiveness of fungicides. *Mol. Plant Microbe Interact.* 14:1114–24
 34. Friedrich L, Lawton K, Reuss W, Masner P, Specker N, et al. 1996. A benzothiadiazole induces systemic acquired resistance in tobacco. *Plant J.* 10:61–70
 35. Gaffney T, Friedrich L, Vernooij B, Negrotto D, Nye G, et al. 1993. Requirement of salicylic acid for the induction of systemic acquired resistance. *Science* 261:754–56
 36. Glazebrook J, Chen W, Estes B, Chang H-S, Nawrath C, et al. 2003. Topology of the network integrating salicylate and jasmonate signal transduction derived from global expression phenotyping. *Plant J.* 34:217–28
 37. Glazebrook J, Rogers EE, Ausubel FM. 1996. Isolation of *Arabidopsis* mutants with enhanced disease susceptibility by direct screening. *Genetics* 143:973–82
 38. Görlach J, Volrath S, Knauf-Beiter G, Hengy G, Beckhove U, et al. 1996. Benzothiadiazole, a novel class of inducers of systemic acquired resistance, activates gene expression and disease resistance in wheat. *Plant Cell* 8:629–43
 39. Guedes MEM, Richmond S, Kuc J. 1980. Induced systemic resistance to anthracnose in cucumber as influenced by the

- location of the inducer inoculation with *Colletotrichum lagenarium* and the onset of flowering and fruiting. *Physiol. Plant Pathol.* 17:229–33
40. Hammond-Kosack KE, Jones JDG. 1996. Resistance gene-dependent plant defense responses. *Plant Cell* 8:1773–91
 41. Heil M. 2002. Ecological costs of induced resistance. *Curr. Opin. Plant Biol.* 5:345–50
 42. Heil M, Baldwin IT. 2002. Fitness costs of induced resistance: emerging experimental support for a slippery concept. *Trends Plant Sci.* 7:61–67
 43. Heil M, Bostock RM. 2002. Induced systemic resistance (ISR) against pathogens in the context of induced plant defences. *Ann. Bot.* 89:503–12
 44. Heil M, Hilpert A, Kaiser W, Linsenmair KE. 2000. Reduced growth and seed set following chemical induction of pathogen defence: Does systemic acquired resistance (SAR) incur allocation costs? *J. Ecol.* 88:645–54
 45. Jakoby M, Weisshaar B, Dröge-Laser W, Vicente-Carbajosa J, Tiedemann J, et al. 2002. bZIP transcription factors in *Arabidopsis*. *Trends Plant Sci.* 7:106–11
 46. Jenns AE, Kuc J. 1979. Graft transmission of systemic resistance of cucumber to anthracnose induced by *Colletotrichum lagenarium* and tobacco necrosis virus. *Phytopathology* 7:753–56
 47. Jirage D, Tootle TL, Reuber TL, Frost LN, Feys BJ, et al. 1999. *Arabidopsis thaliana* *PAD4* encodes a lipase-like gene that is important for salicylic acid signaling. *Proc. Natl. Acad. Sci. USA* 96:13583–88
 48. Johnson C, Boden E, Arias J. 2003. Salicylic acid and NPR1 induce the recruitment of *trans*-activating TGA factors to a defense gene promoter in *Arabidopsis*. *Plant Cell* 15:1846–58
 49. Katagiri F, Lam E, Chua N-H. 1989. Two tobacco DNA-binding proteins with homology to the nuclear factor CREB. *Nature* 340:727–30
 50. Kiefer IW, Slusarenko AJ. 2003. The pattern of systemic acquired resistance induction within the *Arabidopsis* rosette in relation to the pattern of translocation. *Plant Physiol.* 132:840–47
 51. Kim HS, Delaney TP. 2002. Over-expression of *TGA5*, which encodes a bZIP transcription factor that interacts with NIM1/NPR1, confers SAR-independent resistance in *Arabidopsis thaliana* to *Peronospora parasitica*. *Plant J.* 32:151–63
 52. Kinkema M, Fan W, Dong X. 2000. Nuclear localization of NPR1 is required for activation of *PR* gene expression. *Plant Cell* 12:2339–50
 53. Kumar D, Klessig DF. 2003. High-affinity salicylic acid-binding protein 2 is required for plant innate immunity and has salicylic acid-stimulated lipase activity. *Proc. Natl. Acad. Sci. USA* 100:16101–6
 54. Kunkel BN, Brooks DM. 2002. Cross talk between signaling pathways in pathogen defense. *Curr. Opin. Plant Biol.* 5:325–31
 55. Lawton KA, Friedrich L, Hunt M, Weymann K, Delaney T, et al. 1996. Benzothiadiazole induces disease resistance in *Arabidopsis* by activation of the systemic acquired resistance signal transduction pathway. *Plant J.* 10:71–82
 56. Lebel E, Heifetz P, Thorne L, Uknes S, Ryals J, Ward E. 1998. Functional analysis of regulatory sequences controlling *PR-1* gene expression in *Arabidopsis*. *Plant J.* 16:223–33
 57. Lee H-I, León J, Raskin I. 1995. Biosynthesis and metabolism of salicylic acid. *Proc. Natl. Acad. Sci. USA* 92:4076–79
 58. León J, Lawton MA, Raskin I. 1995. Hydrogen peroxide stimulates salicylic acid biosynthesis in tobacco. *Plant Physiol.* 108:1673–78
 59. Li X, Song Y, Century K, Straight S, Ronald P, et al. 2001. A fast neutron deletion mutagenesis-based reverse genetics system for plants. *Plant J.* 27:235–42
 60. Li X, Zhang Y, Clarke JD, Li Y, Dong X. 1999. Identification and cloning of a negative regulator of systemic acquired

- resistance, SNI1, through a screen for suppressors of *npr1-1*. *Cell* 98:329–39
61. Liu Y, Schiff M, Marathe R, Dinesh-Kumar SP. 2002. Tobacco *Rar1*, *EDS1* and *NPR1/NIMI* like genes are required for N-mediated resistance to tobacco mosaic virus. *Plant J.* 30:415–29
 62. Mahalingam R, Gomez-Buitrago A, Eckardt N, Shah N, Guevara-Garcia A, et al. 2003. Characterizing the stress/defense transcriptome of *Arabidopsis*. *Genome Biol.* 4:R20
 63. Malamy J, Carr JP, Klessig DF, Raskin I. 1990. Salicylic acid: a likely endogenous signal in the resistance response of tobacco to viral infection. *Science* 250:1002–4
 64. Maldonado AM, Doerner P, Dixon RA, Lamb CJ, Cameron RK. 2002. A putative lipid transfer protein involved in systemic resistance signalling in *Arabidopsis*. *Nature* 419:399–403
 65. Maleck K, Levine A, Eulgem T, Morgan A, Schmid J, et al. 2000. The transcriptome of *Arabidopsis thaliana* during systemic acquired resistance. *Nat. Genet.* 26:403–10
 66. Mauch F, Mauch-Mani B, Gaille C, Kull B, Haas D, Reimann C. 2001. Manipulation of salicylate content in *Arabidopsis thaliana* by the expression of an engineered bacterial salicylate synthase. *Plant J.* 25:67–77
 67. Mayda E, Mauch-Mani B, Vera P. 2000. *Arabidopsis dth9* mutation identifies a gene involved in regulating disease susceptibility without affecting salicylic acid-dependent responses. *Plant Cell* 12:2119–28
 68. Métraux J-P. 2002. Recent breakthroughs in the study of salicylic acid biosynthesis. *Trends Plant Sci.* 7:332–34
 69. Métraux J-P, Ahl-Goy P, Staub T, Speich J, Steinemann A, et al. 1991. Induced resistance in cucumber in response to 2,6-dichloroisonicotinic acid and pathogens. In *Advances in Molecular Genetics of Plant-Microbe Interactions*, ed. H Henneke, DPS Verma, pp. 432–39. Dordrecht, The Netherlands: Kluwer
 70. Métraux JP, Signer H, Ryals J, Ward E, Wyss-Benz M, et al. 1990. Increase in salicylic acid at the onset of systemic acquired resistance in cucumber. *Science* 250:1004–6
 71. Meuwly P, Mölders W, Buchala A, Métraux J-P. 1995. Local and systemic biosynthesis of salicylic acid in infected cucumber plants. *Plant Physiol.* 109:1107–14
 72. Milligan SB, Bodeau J, Yaghoobi J, Kaloshian I, Zabel P, Williamson VM. 1998. The root knot nematode resistance gene *Mi* from tomato is a member of the leucine zipper, nucleotide binding, leucine-rich repeat family of plant genes. *Plant Cell* 10:1307–19
 73. Mölders W, Buchala A, Métraux J-P. 1996. Transport of salicylic acid in tobacco necrosis virus-infected cucumber plants. *Plant Physiol.* 112:787–92
 74. Mou Z, Fan W, Dong X. 2003. Inducers of plant systemic acquired resistance regulate NPR1 function through redox changes. *Cell* 113:935–44
 75. Nandi A, Welti R, Shah J. 2004. The *Arabidopsis thaliana* dihydroxyacetone phosphate reductase gene *SUPPRESSOR OF FATTY ACID DESATURASE DEFICIENCY 1* is required for glycerolipid metabolism and for the activation of systemic acquired resistance. *Plant Cell* 16:465–77
 76. Nawrath C, Heck S, Parinthewong N, Métraux J-P. 2002. EDS5, an essential component of salicylic acid-dependent signaling for disease resistance in *Arabidopsis*, is a member of the MATE transporter family. *Plant Cell* 14:275–86
 77. Nawrath C, Métraux J-P. 1999. Salicylic acid induction-deficient mutants of *Arabidopsis* express *PR-2* and *PR-5* and accumulate high levels of camalexin after pathogen inoculation. *Plant Cell* 11:1393–404
 78. Neuenschwander U, Vernooij B, Friedrich

- L, Uknes S, Kessmann H, Ryals J. 1995. Is hydrogen peroxide a second messenger of salicylic acid in systemic acquired resistance? *Plant J.* 8:227–33
79. Niggeweg R, Thurow C, Kegler C, Gatz C. 2000. Tobacco transcription factor TGA2.2 is the main component of *as-1*-binding factor ASF-1 and is involved in salicylic acid- and auxin-inducible expression of *as-1*-containing target promoters. *J. Biol. Chem.* 275:19897–905
80. Niggeweg R, Thurow C, Weigel R, Pfitzner U, Gatz C. 2000. Tobacco TGA factors differ with respect to interaction with NPR1, activation potential and DNA-binding properties. *Plant Mol. Biol.* 42:775–88
81. Parker JE, Holub EB, Frost LN, Falk A, Gunn ND, Daniels MJ. 1996. Characterization of *eds1*, a mutation in Arabidopsis suppressing resistance to *Peronospora parasitica* specified by several different *RPP* genes. *Plant Cell* 8:2033–46
82. Pieterse CMJ, van Wees SCM, Hoffland E, van Pelt JA, van Loon LC. 1996. Systemic resistance in Arabidopsis induced by biocontrol bacteria is independent of salicylic acid accumulation and pathogenesis-related gene expression. *Plant Cell* 8:1225–37
83. Pieterse CMJ, van Wees SCM, van Pelt JA, Knoester M, Laan R, et al. 1998. A novel signaling pathway controlling induced systemic resistance in Arabidopsis. *Plant Cell* 10:1571–80
84. Pontier D, Miao Z-H, Lam E. 2001. Transdominant suppression of plant TGA factors reveals their negative and positive roles in plant defense responses. *Plant J.* 27:529–38
85. Rasmussen JB, Hammerschmidt R, Zook MN. 1991. Systemic induction of salicylic acid accumulation in cucumber after inoculation with *Pseudomonas syringae* pv. *syringae*. *Plant Physiol.* 97:1342–47
86. Ross AF. 1961. Systemic acquired resistance induced by localized virus infections in plants. *Virology* 14:340–58
87. Ross AF. 1966. Systemic effects of local lesion formation. In *Viruses of Plants*, ed. ABR Beemster, J Dijkstra, pp. 127–50. Amsterdam: North-Holland
88. Rossi M, Goggin FL, Milligan SB, Kaloshian I, Ullman DE, Williamson VM. 1998. The nematode resistance gene *Mi* of tomato confers resistance against the potato aphid. *Proc. Natl. Acad. Sci. USA* 95:9750–54
89. Ryals J, Lawton KA, Delaney TP, Friedrich L, Kessmann H, et al. 1995. Signal transduction in systemic acquired resistance. *Proc. Natl. Acad. Sci. USA* 92:4202–5
90. Ryals J, Weymann K, Lawton K, Friedrich L, Ellis D, et al. 1997. The Arabidopsis *NIM1* protein shows homology to the mammalian transcription factor inhibitor $\text{I}\kappa\text{B}$. *Plant Cell* 9:425–39
91. Ryals JA, Neuenschwander UH, Willits MG, Molina A, Steiner H-Y, Hunt MD. 1996. Systemic acquired resistance. *Plant Cell* 8:1809–19
92. Schenk PM, Kazan K, Wilson I, Anderson JP, Richmond T, et al. 2000. Coordinated plant defense responses in Arabidopsis revealed by microarray analysis. *Proc. Natl. Acad. Sci. USA* 97:11655–60
93. Shah J, Klessig DF. 1999. Salicylic acid: signal perception and transduction. In *Biochemistry and Molecular Biology of Plant Hormones*, ed. PPJ Hooykaas, MA Hall, KR Libbenga, pp. 513–41. London: Elsevier
94. Shah J, Tsui F, Klessig DF. 1997. Characterization of a salicylic acid-insensitive mutant (*sai1*) of *Arabidopsis thaliana* identified in a selective screen utilizing the SA-inducible expression of the *tms2* gene. *Mol. Plant Microbe. Interact.* 10:69–78
95. Shirano Y, Kachroo P, Shah J, Klessig DF. 2002. A gain-of-function mutation in an Arabidopsis toll interleukin 1 receptor–nucleotide binding site–leucine-rich repeat type *R* gene triggers defense responses and results in enhanced disease resistance. *Plant Cell* 14:3149–62

96. Shirasu K, Nakajima H, Rajasekhar VK, Dixon RA, Lamb C. 1997. Salicylic acid potentiates an agonist-dependent gain control that amplifies pathogen signals in the activation of defense mechanisms. *Plant Cell* 9:261–70
97. Shulaev V, León J, Raskin I. 1995. Is salicylic acid a translocated signal of systemic acquired resistance in tobacco? *Plant Cell* 7:1691–701
98. Shulaev V, Silverman P, Raskin I. 1997. Airborne signalling by methyl salicylate in plant pathogen resistance. *Nature* 385:718–21
99. Silva H, Yoshioka K, Dooner HK, Klessig DF. 1999. Characterization of a new *Arabidopsis* mutant exhibiting enhanced disease resistance. *Mol. Plant Microbe Interact.* 12:1053–63
100. Silverman P, Seskar M, Kanter D, Schweizer P, Métraux J-P, Raskin I. 1995. Salicylic acid in rice (biosynthesis, conjugation, and possible role). *Plant Physiol.* 108:633–39
101. Spoel SH, Koornneef A, Claessens SMC, Korzelius JP, Van Pelt JA, et al. 2003. NPR1 modulates cross-talk between salicylate- and jasmonate-dependent defense pathways through a novel function in the cytosol. *Plant Cell* 15:760–70
102. Sticher L, Mauch-Mani B, Métraux JP. 1997. Systemic acquired resistance. *Annu. Rev. Phytopathol.* 35:235–70
103. Tenhaken R, Rübel C. 1997. Salicylic acid is needed in hypersensitive cell death in soybean but does not act as a catalase inhibitor. *Plant Physiol.* 115:291–98
104. Ton J, Davison S, van Wees SCM, Van Loon LC, Pieterse CMJ. 2001. The *Arabidopsis* *ISR1* locus controlling Rhizobacteria-mediated induced systemic resistance is involved in ethylene signaling. *Plant Physiol.* 125:652–61
105. Ton J, De Vos M, Robben C, Buchala A, Métraux J-P, et al. 2002. Characterization of *Arabidopsis* enhanced disease susceptibility mutants that are affected in systemically induced resistance. *Plant J.* 29:11–21
106. Ton J, Van Pelt JA, Van Loon LC, Pieterse CMJ. 2002. Differential effectiveness of salicylate-dependent and jasmonate/ethylene-dependent induced resistance in *Arabidopsis*. *Mol. Plant-Microbe Interact.* 15:27–34
107. van Loon LC, Bakker PAHM, Pieterse CMJ. 1998. Systemic resistance induced by rhizosphere bacteria. *Annu. Rev. Phytopathol.* 36:453–83
108. van Loon LC, van Kammen A. 1970. Polyacrylamide disc electrophoresis of the soluble leaf proteins from *Nicotiana tabacum* var. ‘Samsun’ and ‘Samsun NN’. II. Changes in protein constitution after infection with tobacco mosaic virus. *Virology* 40:199–211
109. Van Loon LC, Van Strien EA. 1999. The families of pathogenesis-related proteins, their activities, and comparative analysis of PR-1 type proteins. *Physiol. Mol. Plant Pathol.* 55:85–97
110. van Wees SCM, de Swart EAM, van Pelt JA, van Loon LC, Pieterse CMJ. 2000. Enhancement of induced disease resistance by simultaneous activation of salicylate- and jasmonate-dependent defense pathways in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* 97:8711–16
111. Vanacker H, Lu H, Rate DN, Greenberg JT. 2001. A role for salicylic acid and NPR1 in regulating cell growth in *Arabidopsis*. *Plant J.* 28:209–16
112. Verberne MC, Verpoorte R, Bol JF, Mercado-Blanco J, Linthorst HJM. 2000. Overproduction of salicylic acid in plants by bacterial transgenes enhances pathogen resistance. *Nat. Biotechnol.* 18:779–83
113. Vernooij B, Friedrich L, Morse A, Reist R, Kolditz-Jawhar R, et al. 1994. Salicylic acid is not the translocated signal responsible for inducing systemic acquired resistance but is required in signal transduction. *Plant Cell* 6:959–65
114. Ward ER, Uknes SJ, Williams SC,

- Dincher SS, Wiederhold DL, et al. 1991. Coordinate gene activity in response to agents that induce systemic acquired resistance. *Plant Cell* 3:1085–94
115. Weigel RR, Bäuscher C, Pfitzner AJP, Pfitzner UM. 2001. NIMIN-1, NIMIN-2 and NIMIN-3, members of a novel family of proteins from *Arabidopsis* that interact with NPR1/NIM1, a key regulator of systemic acquired resistance in plants. *Plant Mol. Biol.* 46:143–60
116. White RF. 1979. Acetylsalicylic acid (aspirin) induces resistance to tobacco mosaic virus in tobacco. *Virology* 99:410–12
117. Wildermuth MC, Dewdney J, Wu G, Ausubel FM. 2001. Isochorismate synthase is required to synthesize salicylic acid for plant defence. *Nature* 414:562–65
118. Xiang C, Miao Z, Lam E. 1997. DNA-binding properties, genomic organization and expression pattern of *TGA6*, a new member of the *TGA* family of bZIP transcription factors in *Arabidopsis thaliana*. *Plant Mol. Biol.* 34:403–15
119. Yu D, Chen C, Chen Z. 2001. Evidence for an important role of WRKY DNA binding proteins in the regulation of *NPR1* gene expression. *Plant Cell* 13:1527–39
120. Yu D, Liu Y, Fan B, Klessig DF, Chen Z. 1997. Is the high basal level of salicylic acid important for disease resistance in potato? *Plant Physiol.* 115:343–49
121. Zhang Y, Fan W, Kinkema M, Li X, Dong X. 1999. Interaction of NPR1 with basic leucine zipper protein transcription factors that bind sequences required for salicylic acid induction of the *PR-1* gene. *Proc. Natl. Acad. Sci. USA* 96:6523–28
122. Zhang Y, Goritschnig S, Dong X, Li X. 2003. A gain-of-function mutation in a plant disease resistance gene leads to constitutive activation of downstream signal transduction pathways in *suppressor of npr1-1, constitutive 1*. *Plant Cell* 15:2636–46
123. Zhang Y, Tessaro MJ, Lassner M, Li X. 2003. Knockout analysis of Arabidopsis transcription factors *TGA2*, *TGA5*, and *TGA6* reveals their redundant and essential roles in systemic acquired resistance. *Plant Cell* 15:2647–53
124. Zhou J-M, Trifa Y, Silva H, Pontier D, Lam E, et al. 2000. NPR1 differentially interacts with members of the TGA/OBF family of transcription factors that bind an element of the *PR-1* gene required for induction by salicylic acid. *Mol. Plant Microbe Interact.* 13:191–202
125. Zhou N, Tootle TL, Tsui F, Klessig DF, Glazebrook J. 1998. *PAD4* functions upstream from salicylic acid to control defense responses in Arabidopsis. *Plant Cell* 10:1021–30



CONTENTS

FRONTISPIECE, <i>Anne K. Vidaver</i>	x
THE ACCIDENTAL PLANT PATHOLOGIST, <i>Anne K. Vidaver</i>	1
TOBACCO MOSAIC VIRUS: A MODEL SYSTEM FOR PLANT BIOLOGY, <i>Karen-Beth G. Scholthof</i>	13
ASSESSMENT AND MANAGEMENT OF SOIL MICROBIAL COMMUNITY STRUCTURE FOR DISEASE SUPPRESSION, <i>Mark Mazzola</i>	35
ANALYSIS OF DISEASE PROGRESS AS A BASIS FOR EVALUATING DISEASE MANAGEMENT PRACTICES, <i>M.J. Jeger</i>	61
EVOLUTION OF PLANT PARASITISM AMONG NEMATODES, <i>J.G. Baldwin, S.A. Nadler, and B.J. Adams</i>	83
LESSONS LEARNED FROM THE GENOME ANALYSIS OF <i>RALSTONIA SOLANACEARUM</i> , <i>Stéphane Genin and Christian Boucher</i>	107
MANAGEMENT AND RESISTANCE IN WHEAT AND BARLEY TO FUSARIUM HEAD BLIGHT, <i>Guihua Bai and Gregory Shaner</i>	135
COMPARATIVE GENOMICS ANALYSES OF CITRUS-ASSOCIATED BACTERIA, <i>Leandro M. Moreira, Robson F. de Souza, Nalvo F. Almeida Jr., João C. Setubal, Julio Cezar F. Oliveira, Luiz R. Furlan, Jesus A. Ferro, and Ana C.R. da Silva</i>	163
SYSTEMIC ACQUIRED RESISTANCE, <i>W.E. Durrant and X. Dong</i>	185
MOLECULAR ASPECTS OF PLANT VIRUS TRANSMISSION BY OLPIDIUM AND PLASMIDIOPHORID VECTORS, <i>D'Ann Rochon, Kishore Kakani, Marjorie Robbins, and Ron Reade</i>	211
MICROBIAL DIVERSITY IN SOIL: SELECTION OF MICROBIAL POPULATIONS BY PLANT AND SOIL TYPE AND IMPLICATIONS FOR DISEASE SUPPRESSIVENESS, <i>P. Garbeva, J.A. van Veen, and J.D. van Elsas</i>	243
MICROBIAL DYNAMICS AND INTERACTIONS IN THE SPERMOSPHERE, <i>Eric B. Nelson</i>	271
BIOLOGICAL CONTROL OF CHESTNUT BLIGHT WITH HYPOVIRULENCE: A CRITICAL ANALYSIS, <i>Michael G. Milgroom and Paolo Cortesi</i>	311
INTEGRATED APPROACHES FOR DETECTION OF PLANT PATHOGENIC BACTERIA AND DIAGNOSIS OF BACTERIAL DISEASES, <i>Anne M. Alvarez</i>	339

NEMATODE MOLECULAR DIAGNOSTICS: FROM BANDS TO BARCODES, <i>Tom Powers</i>	367
TYPE III SECRETION SYSTEM EFFECTOR PROTEINS: DOUBLE AGENTS IN BACTERIAL DISEASE AND PLANT DEFENSE, <i>Allan Collmer</i> <i>and James R. Alfano</i>	385
PLANT VIRUS SATELLITE AND DEFECTIVE INTERFERING RNAs: NEW PARADIGMS FOR A NEW CENTURY, <i>Anne E. Simon, Marilyn J. Roossinck,</i> <i>and Zoltán Havelda</i>	415
CHEMICAL BIOLOGY OF MULTI-HOST/PATHOGEN INTERACTIONS: CHEMICAL PERCEPTION AND METABOLIC COMPLEMENTATION, <i>Andrew G. Palmer,</i> <i>Rong Gao, Justin Maresh, W. Kaya Erbil, and David G. Lynn</i>	439
INDEX Subject Index	465
ERRATA An online log of corrections to <i>Annual Review of Phytopathology</i> chapters may be found at http://phyto.annualreviews.org/	